

erc

Progenitors of LIGO's Black Holes

The era of GW Astronomy, Paris, June 26-20, 2017



edit background NASA Paresce, Design E. Buunk

Bin
Cosmos

Amsterdam



Manos
Zapartas
PhD



Ylva
Göteborg
PhD



Mathieu
Renzo
PhD




Silvia
Toonen
VENI fellow




6 new
PhD/PD

Interested?



Louise
Edstam
MSc



Walter
Rossem
MSc



Thomas
Dodds
BSc



Rob
Farmer
Postdoc



Eva
Laplace
PhD



Selma E. de Mink



Anton Pannekoek Inst., University of Amsterdam



erc

Progenitors of LIGO's Black Holes

The era of GW Astronomy, Paris, June 26-20, 2017



Ilya Mandel, Rob Izzard, Hugues Sana, Pablo Marchant, Chris Belczynski, Andrew King, Philip Podsiadlowski, Simon Stevenson, Alejandro Vigna-Gómez, Norbert Langer, Fabian Schneider, Stephen Justham, Sung-Chul Yoon, Ori Fox, Schuyler van Dyk, Jose Groh, Nathan Smith, , Carles Badenes, Alex de Koter, Tom Maccarone, Leo Almeida, Tony Piro, Maria Drout, Coen van Neijssel, Abel Schootemeijer, Colin Norman,

VLT-FLAMES Massive Star Consortium (PI: Evans)



edit background NASA Patresce, Design E. Buunk


Selma E. de Mink

Anton Pannekoek Inst., University of Amsterdam

How did they form?



How did they form?



Scope

(A) Stellar Origin Black Holes

- ✓ Gravitational Collapse of (the core of) a massive star
- ✓ Observed (albeit at somewhat lower mass)




(B) Primordial Black Holes


Carr & Hawking (1974), Carr (1975, 1976), Garcia-Bellido et al. (1996), Khlopov (2010), Frampton et al. (2010), Blais et al. (2002), ...

- ✓ Gravitational Collapse of extreme densities at high redshift
- ✓ Still hypothetical ... (so far)
- ✓ Somehow evaded strong constraints by
 - Micro lensing events
 - CMB spectral distortions
- ✓ Constituent of Dark Matter (instead / in addition to more popular but still elusive WIMPs)


Known black holes that have a stellar origin (presumably)



Are LIGO's black holes of stellar origin?




Are LIGO's black holes of stellar origin?



A white silhouette of a person with long, flowing hair, possibly a woman, is positioned on the left side of the slide. The person's arms are raised, and their hair is blowing in the wind, creating a sense of movement. The silhouette is set against a solid grey background.

When did LIGO's
black holes form?



When did the progenitors of LIGO's black holes form?


- 1. Such that they merge within LIGO's reach.
(optimum between: detection horizon & delay time)**
- 2. When the Universe was forming stars at high rate**
- 3. When "conditions" where "right"
(to form binary black holes, e.g. metallicity)**

1.


Birth of progenitors stars



Formation black holes



Coalescence



t_0


Time

Lifetime stellar
progenitors
~ 10 Myrs



In-spiral due to
Gravitational Waves
Model dependent: Typically: ~ Gyrs

Detectable
Signal
~ 0.1 second

Birth of progenitors Birth black holes




Coalescence




You are here


2. Cosmic Star formation history





2. Cosmic Star formation history



Birth of progenitors Birth black holes



Coalescence



dark
ages

First
Stars

time



You are
here


A white silhouette of a person with long, flowing hair, possibly a dancer or a person in motion, set against a grey background.

Which stars make black holes?

A blue arrow pointing downwards and to the right, containing the text "Small print".


Small print

Disclaimer: we don't know.
Any assumptions you make are at your own risk



Final fate of stars

How hard can it be ...? We all know this ...



NASA outreach material.

Stellar Evolution

How hard can it be ... ?

Four coupled differential equations

I. Conservation of Mass

$$\frac{dr}{dM_r} = \frac{1}{4\pi r^2 \rho}$$

“Heroic Struggle against gravity”

$$\frac{dP}{dM_r} = -\frac{GM_r}{4\pi r^4}$$

III. Conservation of Energy

$$\frac{dL_r}{dM_r} = \epsilon$$

IV. How is energy transported

$$\frac{dT}{dM_r} = -\frac{1}{16\pi^2 r^4 \lambda \rho} L_r$$

Independent variable

$$M_r.$$

Four primary variables

r, P, L_r and $T,$

Boundary conditions:

$$r = 0, \quad L_r = 0, \quad \text{for } M_r = 0$$

$$P = 0, \quad T = 0, \quad \text{for } M_r = M,$$

Auxiliary equations

ρ : equation of state, $P = P(\rho, T, X_i)$

λ : coefficient of conductivity, $\lambda(\rho, T, X_i)$

ϵ : nuclear fusion rate, $\epsilon(\rho, T, X_i)$

Nuclear reaction network

2 types of simulations

Rapid evolutionary 'prescriptions'


- Computationally cheap → used to make predictions for populations / rates
- StarTrack, binary_c, BSE, ...


Full solutions to stellar evolution equations

- More expensive computationally, (in part due to convergence issues)
- Kippenhahn codes, MESA




Mass loss





Metallicity




Z : Mass fraction of all elements more massive than H & He

$[\text{Fe}/\text{H}]$, $[\text{O}/\text{H}]$: Log_{10} of ratio compared to the sun

Why do stars care about Metallicity

Mass loss by (radiatively driven) stellar winds cares about metals




Mathieu Renzo

Reduced (line driven) winds at lower metallicity

Vink et al. 2000, Mokiem et al. 2005

“Mass Loss”



Challenge 1 wind clumping?

ANTON PANNEKEK
INSTITUT

Inhomogeneities:

$$f_{\text{cl}} \stackrel{\text{def}}{=} \frac{\langle \rho^2 \rangle}{\langle \rho \rangle^2} \neq 1 \Rightarrow \dot{M} \neq 4\pi r^2 \rho v(r)$$


4/



Mathieu
Renzo


Mass loss uncertainties

Renzo et al. (2017)



Mass loss uncertainties


Renzo et al. (2017)



Mathieu Renzo

Mass loss uncertainties


Renzo et al. (2017)




**Mathieu
Renzo**

Challenge 2: Other poorly understood modes of mass loss

Eruptive Mass loss




Stripping by Companion



Need for reduced winds

Belczynski et al. 2010


Astrophysical implications paper



Need for reduced winds


Belczynski et al. 2010

Astrophysical implications paper




A white silhouette of a person with long, flowing hair, possibly a dancer or a person in motion, set against a solid grey background. The figure is positioned on the left side of the frame, with arms raised and hair blowing outwards.

Which (single) stars make black holes



How massive single stars end their lives? - 1


As we thought about this
a decade ago



Maybe not that simple:


“Islands” of black hole formation.

Current discussion



Maybe not that simple


O. Pejcha¹ and T. A. Thompson²



A white silhouette of a person's head and shoulders, looking upwards with their arms raised, set against a grey background.

What do we know about the progenitors

Closest Peak at their Progenitors



Large Magellanic Cloud
 $Z / Z_0 \approx 1/2$

Small Magellanic Cloud
 $Z / Z_0 \approx 1/5$


Black Holes in the making: Tarantula Nebula



VLT-FLAMES ([Evans](#), Sana), HST ([Sabbi](#), Lennon, Crowther), Chandra ([Townsend](#))

Black Holes in the making: Tarantula Nebula

Hertzsprung-Russell diagram



- Densest sampling of the upper stellar mass function available (up to 200M $_{\odot}$)
- **Excess of massive stars** above 30M $_{\odot}$ ($32 \pm 12\%$ more than predicted by Salpeter IMF in mass range 15–200M $_{\odot}$)
- **Flatter IMF slope:** $1.90^{+0.37}_{-0.26}$ (cf. Salpeter = 2.35)

1. Masses





1. Masses

Nine "Monster" stars

Crowther+10,+15


Current masses:
~ 100 - 250 $M_{\odot}$



3 Binaries everywhere ...





This is not exotic




cf. Abt+78, Kobulnicky+Fryer07, Mason+09,
Chini+12, Kobulnicky+14, Sana+12, Sana+13,
Dunstall+15, Moe+16, Almeida+17, ...

How do we know?




How do we know?





www.eso.org


VFTS 352
Orbit: 1.1 Day



Will they merge?
Will they mix and shrink?

Almeida, Sana, de Mink et al. (2015),
dM+09, Mandel+dM16, Marchant+16,17
Image Credit:ESO/L. Calçada


How do we know?



Evolution of a single star

MESA (Paxton+11,13,15): $M_1 = 15 M_\odot$

Götberg et al. (2017a)




Ylva
Götberg

Evolution of a very typical donor star

He Giant

$M_1 = 15 M_\odot$ $q = 0.8$ *Orbital Period: 10 d*

Götberg et al. (2017a)




cf. Kippenhahn, Podsiadlowski, Langer, Wellstein, Pols, Yoon, Claeys, Eldridge, Bersten ...

Donor star

MESA: $M_1 = 15 M_\odot$ $q = 0.8$

Götberg et al. (in prep)




cf. Kippenhahn, Podsiadlowski, Langer, Wellstein, Pols, Yoon, Claeys, Eldridge, Bersten ...

Donor star

MESA: $M_1 = 15 M_\odot$ $q = 0.8$

Götberg et al. (in prep)




cf. Kippenhahn, Podsiadlowski, Langer, Wellstein, Pols, Yoon, Claeys, Eldridge, Bersten ...

Donor star

MESA: $M_1 = 15 M_\odot$ $q = 0.8$

Götberg et al. (in prep)




cf. Kippenhahn, Podsiadlowski, Langer, Wellstein, Pols, Yoon, Claeys, Eldridge, Bersten ...

Donor star

MESA: $M_1 = 15 M_\odot$ $q = 0.8$

Götberg et al. (in prep)






cf. Kippenhahn, Podsiadlowski, Langer, Wellstein, Pols, Yoon, Claeys, Eldridge, Bersten ...

Animation of the life of a typical massive Star



www.eso.org

Majority (75-98%) disrupt when the first star explodes



Runaway stars
(or “walkaway stars”)

Zapartas, dM +17b
Renzo & dM (in prep)

Cf. Zwicky57, Blaauw61, Boersma61, Brandt+Podsiadlovski95,
Kalogera96, Fryer+98, Dray+05, Eldridge+11, Sana+prep




Manos
Zapartas




Mathieu
Renzo

Some binaries survive




Some binary stars stay together forever



A white silhouette of a person's head and shoulders, looking upwards with their arms raised, set against a light gray background.

**Why is it so hard
to form binary black holes?**



A light gray silhouette of a man and a woman in a dynamic dance pose. The man is on the left, leaning back with his arms raised, and the woman is on the right, leaning forward with her arms raised. The background is white, and the text is centered over the image.

Two Main Challenges for all progenitor scenarios

Progenitors




***0. Need to overcome their
“Relationship issues”***

1. “Separation Challenge”


2. “Mass Challenge”

Separation



Using Peters '64

Separation



$R_o \sim 7e10$ cm

Using Peters '64

Separation

1. “Separation Challenge”

*How to get Black holes close enough
to coalesce in a Hubble time?*

$10 R$




2. “Mass Challenge”

*How to avoid
excessive Mass loss?*

Two types of Formation Channels

1. Evolutionary
formation channels



2. Dynamical
formation channels



Stellar Density




Two types of Formation Channels

1. Evolutionary formation channels

- ✓ **Classical channel**
(involving common envelope or other forms of mass transfer)
- ✓ **Chemically Homogeneous Channel**
(mixing processes in near contact binaries)
- ✓ Population III binaries
- ✓ Stable mass transfer ...


2. Dynamical formation channels


- ✓ Chaotic Dynamics in **dense Star Clusters** or Nuclear star clusters
- ✓ "" in Nuclear star clusters in center of galaxies
- ✓ "" Near a supermassive black hole, possibly with **AGN discs**
- ✓ Dynamics in lower mass clusters
- ✓ Resonances in **Triple systems**
- ✓ ...

A white silhouette of a person with long, flowing hair, positioned on the left side of the slide. The person's head is tilted back, and their arms are raised, with hands near their head.

Chemically Homogeneous Channel




deMink+08, +09,
Mandel+deMink16, deMink+Mandel16
Marchant+16, 17








DeMink++08, 09,
Mandel+deMink16, deMink+Mandel16
Marchant+16, 17

Standard Evolution




Time
↓

Chemically Homogeneous




Do the Binary BHs merge?



✓ Yes ..
within a Hubble time


Mandel & De Mink (2016) cf. Marchant et al. (2016)

De Mink & Mandel (2016)



What about the spins?



Low spins from metallicities $Z_{\odot} / 10$



Marchant + 2016,
De Mink+ in prep, Marchant in prep.


Side products: what if only one star evolves chemically homogeneous

ULX: Marchant + 2017, Pair Instability Supernova: Du Boisson + to be submitted



A white silhouette of a person with long, flowing hair, positioned on the left side of the slide. The person's head is tilted back, and their arms are raised, with hands near their head. The background is a solid light gray.

Wrap up



Main challenges that (Single) Star evolution faces

- How much mass do stars lose?
- How much mixing occurs (how big are their cores)?
- What decides whether a star forms a NS or BH?
- What are the kicks?

Recent Insight from (EM) observations

- Binarity very common
- IMF extends above 150 M_{sun}
- Possibly evidence for excess of massive stars

Formation scenarios:


- Explanations for the Mass and Separation Challenge

Homogeneous evolution scenario

Making-of


(i)

“Classical” Channel




(ii)

Dynamical interactions



(iii)

“Homogeneous evolution”
in tidally distorted binaries



... Positions opening in Amsterdam

